

Signal Restoration with the Parametric Wiener Filter

Ken Castleman, 3/24/94
Section 16.2.4

pwf_1D.MCD

$$N_p := 20 \quad u := 0, \frac{0.5}{N_p} \dots 0.5$$

The Gaussian: $G(\mu, \sigma, x) \equiv \text{if} \left[|x - \mu| < 3 \cdot \sigma, e^{-\left[\frac{(x-\mu)^2}{(2 \cdot \sigma \cdot \sigma)} \right]}, 0 \right]$

The signal, $f(x)$, has been smoothed by $H(u)$ and corrupted by additive white noise of power N .



$$\mu \equiv 0 \quad \sigma \equiv 0.2 \quad N \equiv 0.1 \quad k \equiv 100$$

The smoothing MTF: $H(u) := G(\mu, \sigma, u)$

The power spectra: $P_f(u) := \frac{2}{1 + k \cdot u^2}$

$P_n(u) := N$

The transfer function:
$$K(u, \gamma) := \frac{H(u)}{(H(u))^2 + \gamma \cdot \left(\frac{P_n(u)}{P_f(u)} \right)}$$



Parametric Wiener Filter Transfer Function

Note that $\gamma = 0$ is straight deconvolution, $\gamma = 1$ is Wiener deconvolution, and $0 < \gamma < 1$ is everything in between. In this case the signal dominates the noise over almost the entire spectrum, but the blurring is rather severe. $\gamma = 0.02$ gives a pretty good looking restoration MTF. Wiener deconvolution is usually too mild.